arXiv:1401.2432v1 [nucl-ex] 10 Jan 2014

d+Au Hadron Correlation Measurements at PHENIX

Anne M. Sickles, for the PHENIX Collaboration

Physics Department, Brookhaven National Laboratory, Upton NY

Abstract

In these proceedings, we discuss recent results from d+Au collisions in PHENIX ridge related measurements and their possible hydrodynamic origin. We present the v_2 at midrapidity and measurements of the pseudorapidity dependence of the ridge, distinguishing between the d-going and Au-going directions. We investigate the possible geometrical origin by comparing v_2 in d+Au to that in p+Pb, Au+Au and Pb+Pb collisions. Future plans to clarify the role of geometry in small collision systems at RHIC are discussed.

Keywords:

In heavy ion collisions at both RHIC and the LHC the properties of the created matter are understood to be described by hydrodynamics with a very small shear viscosity to entropy density ratio, η/s [1]. The value of η/s is constrained via measurements of Fourier coefficients of the azimuthal distribution of particles (v_N where N is the order of the Fourier coefficient). Measurements of the particle pair correlations in heavy ion collisions have been well described by products of the same v_N at the appropriate p_T [2, 3]. One prominent feature of these correlation functions is the so-called *ridge*, a long range in pseudorapidity, small $\Delta\phi$ correlation resulting from the sum of positive v_N from hydrodynamic flow [4, 2].

Surprisingly, a similar long range correlation was observed in very high multiplicity p+p collisions at the LHC [5] where a hydrodynamical system was not generally expected to be created. Recently, at the LHC a double ridge structure, with long range correlations at $\Delta\phi = 0$ and π has been observed also in p+Pb collisions at 5.02 TeV [6, 7]. This feature can be largely described by a $\cos 2\Delta\phi$ modulation. Extractions of v_2 yielded values with a

Email address: annebnl.gov (Anne M. Sickles, for the PHENIX Collaboration)



Figure 1: v_2 as a function of p_T for midrapidity hadrons in the 5% most central d+Au collisions. Also shown on the plot are hydrodynamic calculations from Refs. [9, 10, 11, 12]. Figure is from Ref. [13].

similar magnitude to those in heavy ion collisions [6, 7]. Obviously, such similar behavior between A+A and p(d)+A collisions is suggestive of a similar physical origin. This was quite surprising considering the small size of the overlap region between the two nuclei in p(d)+A collisions. The observation of a double ridge structure, of course, does not prove hydrodynamic behavior; in fact, similar effects were expected within the Color Glass Condensate model [8]. Therefore, experimentally, it is of great interest to investigate the nature of the correlations observed in p(d)+A and to compare measurements in p(d)+A to what is known from A+A collisions. Approximately 1.6 billion d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV was taken in 2008. Here we report on measurements by the PHENIX collaboration using that data.

1. Midrapidty Correlations & v_2

PHENIX has measured the v_2 of charged hadrons at midrapidity in central d+Au collisions using two-particle correlations. Correlations from jets are



Figure 2: v_2/ε_2 as a function of the midrapidity charged particle multiplicity in d+Au, p+Pb, Au+Au and Pb+Pb collisions. The ε_2 values are calculated within a Glauber Monte Carlo. Default ε_2 values are obtained representing the nucleons as point-like centers. Also shown are results with the nucleons represented as Gaussians with $\sigma = 0.4$ fm (red dashed line) and solid disks with radius of 1 fm (green dotted line). Figure is from Ref. [13].

removed under the assumption that the conditional yields are unmodified from peripheral d+Au events. Resulting v_2 values as a function of p_T are shown in Fig. 1 [13]. The v_2 rises with p_T reaching a maximal value of about 15%. Also shown on the plot are hydrodynamic calculations from three groups [9, 10, 11, 12]. All three calculations agree well with the data. Refs. [9, 10, 11] use $\eta/s = 0.08$, while the calculation in Ref. [12] is for ideal hydrodynamics.

In order to further investigate the possible relationship between the geometry of the collision system and the observed v_2 we compare the v_2 values (at $p_T \approx 1.4$ GeV/c) scaled by estimates of the initial state second order eccentricity, ε_2. The v_2/ε_2 values are shown as a function of $dN_{ch}/d\eta$ at mid-rapidity in Fig. 2 for d+Au, p+Pb, Au+Au and Pb+Pb collisions at both RHIC and the LHC. The ε_2 values are from a Glauber Monte Carlo calculation; v_2/ε_2 is consistent between central d+Au and midcentral p+Pb (systems which have a similar $dN_{ch}/d\eta$) despite the factor of 25 difference in collision energy per nucleon pair. The ε_2 value in central d+Au collisions is approximately 50% larger than in in midcentral p+Pb. The v_2/ε_2 ratio rises as a function $dN_{ch}/d\eta$ and follows approximately a common trend between


Au+Au, ε2(point-like centers) (√s=200 GeV)
ε2(Gaussian smearing σ = 0.4)
ε2(disks, R = 1fm)
d+Au, ε2(point-like centers) (√s=200 GeV)
ε2(Gaussian smearing σ = 0.4)
ε2(disks, R = 1fm)
CMS Pb+Pb v2{EP}/ε2part (√s=2.76 TeV)
CMS Pb+Pb v2{2}/ε2{2} (√s=2.76 TeV)
⟨pT⟩ ≈ 1.4 GeV/c
v2 / ε2
1 / S dNch /dη|η=0 (fm-2)

Figure 3: v_2/ε_2 as a function of $\frac{1}{S}\frac{dN_{ch}}{d\eta}$ in d+Au [13], Au+Au [14, 15] and Pb+Pb [16] collisions. The ε_2 and S values are calculated within a Glauber Monte Carlo. Default values represent the nucleons as point-like centers. Also shown for the Au+Au and d+Au points are results with the nucleons represented as Gaussians with $\sigma = 0.4$ fm (red dashed line and red open circle) and solid disks with radius of 1 fm (green dotted line and green open cross).

the four collisions systems.

There are uncertainties within the Glauber Monte Carlo calculation of the spatial eccentricity, ε_2, values. We have investigated the uncertainty due to the modeling of the nucleons within the calculation. The default values take the nucleons as point-like centers and we have also investigated treating the nucleons as disks with a radius of 1 fm and as Gaussians with $\sigma = 0.4$ in both d+Au and Au+Au collisions. In d+Au collisions these variations change ε_2 by approximately 30% at maximum; in Au+Au the effect is much smaller.

In addition to the multiplicity scaling shown in Fig. 2 it is also of interest to investigate the scaling of the same v_2/ε_2 data as a function of the charged particle multiplicity divided by S, a measure of the overlap area between the two nucleons which is defined as $S \equiv 4\pi\sqrt{\sigma_x^2\sigma_y^2 - \sigma_{xy}^2}$ where $\sigma_{xy} \equiv \langle xy\rangle - \langle x\rangle\langle y\rangle$. This scaling has been found to approximately hold between RHIC and the LHC for p_T integrated v_2/ε_2 values [16]. In Fig. 3 v_2/ε_2 is plotted as a function of $\frac{1}{S}\frac{dN_{ch}}{d\eta}$ at $p_T \approx 1.4$ GeV/c. We observe this scaling to hold in d+Au, Au+Au and Pb+Pb collisions. The same three assumptions about the nucleon representation within the Glauber model are used here as well.

In this case, the different nucleon representations affect both the values of ε_2 and S.

2. Rapidity Separated Correlations

In order to investigate whether the correlations seen at midrapidity are long-range in pseudorapidity and, if so, what the pseudorapidity dependence is, we have measured correlations between mid-rapidity hadrons ($|\eta|$ <0.35) and electromagnetic energy in the Au-going and d-going Muon Piston Calorimeters (MPC) [17]. These calorimeters are positioned on either side of the interaction region at $3.1 < |\eta| < 3.7$ (3.9) in the Au-going (d-going) direction. With this large $\Delta\eta$ separation, no same-side jet correlations remain.

In d+Au collisions there is a large asymmetry in the number of particles produced at large pseudorapidity between the *d*-going and Au-going directions. In the most central 20% of the collisions approximately four times as many charged particles are produced in the Au-going direction as in the *d*-going direction [18].

The azimuthal correlations for both the d-going and Au-going MPC with midrapidity tracks are shown in Fig. 4 as a function of centrality. In all cases the dominant feature of the correlation functions is the peak at $\Delta\phi$=π. This peak has contributions from jet correlations and momentum conservation.

In the d-going correlations no near side correlations are seen at any centrality. In the Au-going correlations, peripheral correlations look similar to those observed in the d-going direction. However, in central collisions a positive correlation at $\Delta\phi = 0$ is observed for the 20% most central events. The magnitude of the correlation relative to the peak at $\Delta\phi$=π increases toward more central events. This provides evidence that the v_2 observed at midrapidity is from a long-range pseudorapidity correlation.

3. Future Investigations

While many of the existing measurements in p+Pb and d+Au are suggestive of hydrodynamic behavior in d+Au collisions, further investigations of this surprising effect are certainly in order. No significant v_3 in d+Au collisions was observed [13]. This is consistent with the large v_2 driven by the elongated shape of the deuteron and, as discussed above, the v_3 values extracted from the d+Au data are much smaller than v_2 and consistent with zero. If the geometry of the deuteron causes the observed v_2 values then it



Figure 4: Azimuthal correlations between mid-rapidity hadrons and E_T as measured in the MPC in the d-going (left panels) and Au-going (right panels) direction. The centrality selections are as noted on the plots.

could be possible to induce a large v_3 by using a projectile nucleus with a large ε_3. PHENIX has proposed running ^{3}He+Au, d+Au and p+Au collisions at RHIC in 2015 to directly investigate the relationship between v_N and geometry.

Fig. 5 shows the results of Glauber Monte Carlo calculations of ^{3}He+Au, d+Au and p+Au collisions. The ε_2 value decreases with an increasing number of binary nucleon-nucleon collisions (N_{coll}) in p+Au collisions while for d+Au and ^{3}He+Au collisions it increases up to a maximal value of approximately 0.5. The ε_3 in ^{3}He+Au collisions reaches 0.25 at about 10 collisions and remains approximately constant for all more central collisions. In the 0-5% central d+Au collisions studied here the mean N_{coll} value is about 18, thus the ε_3 value in central ^{3}He+Au collisions will be approximately 40% bigger than in d+Au collisions.

Additionally, in the 2015 run PHENIX will have additional tracking, compared to the measurements presented here, provided by the silicon vertex detectors. This will provides larger $\Delta\eta$ coverage around midrapidity. Thus, it will be possible to determine whether the increased ε_3 in ^{3}He+Au collisions leads to a correspondingly large v_3 as a conclusive test of the role of geometry



Figure 5: (left)An event display of a ^{3}He+Au collision as simulated in a Glauber Monte Carlo. The nucleons are modeled as Gaussians with σ=0.4. The outlines of the nucleons are shown as circles and the color axis shows the Glauber initial energy density. (middle) ε_2 as calculated in a Glauber Monte Carlo as a function of N_{coll} for p+A (black), d+Au (red) and ^{3}He+Au (blue) collisions. (right) ε_3 as a function of N_{coll}. Colors are as in the middle panel.

in generating the v_N observed in p+A and d+A collisions. Recently, hydrodynamic calculations of v_N in p+Au, d+Au and ^{3}He+Au collisions have been done [19] and they provide support for pursing this program.

4. Conclusions

Over the past year, there has been much excitement around the novel effects observed in p+Pb and d+Au collisions. We have presented PHENIX results on the d+Au v_2 at midrapidity and shown new results using our MPC detector which provide the first evidence for long range ($\Delta\eta \approx 3.5$) correlations at small $\Delta\phi$ in d+Au collisions. We have also discussed future plans to constrain the role of geometry in small collision systems by varying the shape of the projectile nucleus in order to significantly vary the ε_3 value of the collision region. The collision system variation possible at RHIC make feature studies in this direction especially exciting.

References

[1] A. Adare, et al., Phys.Rev.Lett. 107 (2011) 252301. arXiv:1105.3928.

[2] K. Aamodt, et al., Phys.Lett. B708 (2012) 249–264. arXiv:1109.2501.

[3] G. Aad, et al., Phys.Rev. C86 (2012) 014907. arXiv:1203.3087.

[4] B. Abelev, et al., Long range rapidity correlations and jet production in high energy nuclear collisions, Phys.Rev. C80 (2009) 064912. arXiv:0909.0191, doi:10.1103/PhysRevC.80.064912.

[5] V. Khachatryan, et al., JHEP 1009 (2010) 091. arXiv:1009.4122.

[6] B. Abelev, et al., Phys.Lett. B719 (2013) 29–41. arXiv:1212.2001.

[7] G. Aad, et al., Phys.Rev.Lett. 110 (2013) 182302. arXiv:1212.5198.

[8] K. Dusling, R. Venugopalan, Phys.Rev. D87 (2013) 054014. arXiv:1211.3701.

[9] P. Bozek, Phys.Rev. C85 (2012) 014911. arXiv:1112.0915.

[10] P. Bozek, private communication.

[11] A. Bzdak, B. Schenke, P. Tribedy, R. VenugopalanarXiv:1304.3403.

[12] G.-Y. Qin, B. MullerarXiv:1306.3439.

[13] A. Adare, et al., Phys.Rev.Lett.arXiv:1303.1794, doi:10.1103/PhysRevLett.111.212301.

[14] S. Adler, et al., Phys.Rev. C71 (2005) 034908. arXiv:nucl-ex/0409015, doi:10.1103/PhysRevC.71.049901,10.1103/PhysRevC.71.034908.

[15] R. A. Lacey, A. Taranenko, R. Wei, N. Ajitanand, J. Alexander, et al., Phys.Rev. C82 (2010) 034910. arXiv:1005.4979, doi:10.1103/PhysRevC.82.034910.

[16] S. Chatrchyan, et al., Phys.Rev. C87 (2013) 014902. arXiv:1204.1409, doi:10.1103/PhysRevC.87.014902.

[17] A. Adare, et al., Phys.Rev.Lett. 107 (2011) 172301. arXiv:1105.5112.

[18] B. Back, et al., Phys.Rev. C72 (2005) 031901. arXiv:nucl-ex/0409021, doi:10.1103/PhysRevC.72.031901.

[19] J. Nagle, A. Adare, S. Beckman, T. Koblesky, J. O. Koop, et al.arXiv:1312.4565.